

# TEST REPORT

8605 SW Creekside Place  
Beaverton, OR 97008  
Phone: 503-466-2445 Fax: 503-466-1636



# D2020 07 07 1079 B

Ordering Provider:  
Getuwell

Samples Received  
07/07/2020

Report Date  
07/07/2020

Samples Collected  
Blood Spot - 07/05/20 08:00

Patient Name: Test Patient  
Patient Phone Number:

Gender	Height	Waist
Male	Unspecified	Unspecified
DOB	Weight	
7/22/1970 (49 yrs)	Unspecified	

TEST NAME	RESULTS   07/05/20	RANGE
Blood Spot		
COVID-19 IgG (Nucleocapsid) §	<b>POSITIVE</b>	Not Applicable
COVID-19 IgG (S1 Spike) §	<b>POSITIVE</b>	Not Applicable

<dl = Less than the detectable limit of the lab. N/A = Not applicable; 1 or more values used in this calculation is less than the detectable limit. H = High. L = Low.

§ ZRT Laboratory COVID-19 IgG antibody serology testing statement

- This test has not been reviewed by the FDA.
- Negative results do not preclude acute SARS-CoV-2 infection. If acute infection is suspected, direct testing for SARS-CoV-2 is necessary.
- Results from antibody testing should not be used to diagnose or exclude acute SARS-CoV-2 infection.
- Positive results may be due to past or present infection with non-SARS-CoV-2 coronavirus strains, such as coronavirus HKU1, NL63, OC43, or 229E.

## Therapies

None

## Lab Comments

Your result for IgG antibodies against the SARS-CoV-2 nucleocapsid protein is positive.

Your result for IgG antibodies against the SARS-CoV-2 S1 spike protein is positive.

Both IgG antibodies are positive, indicating a recent COVID-19 infection. You may or may not have experienced some of the most common symptoms of COVID-19, which include cough, fever, chills, muscle pain, headache, fatigue, sore throat, diarrhea, and loss of taste or smell. Some people don't display any symptoms (asymptomatic). Since our understanding of this disease is still progressing, it is unclear if forming antibodies will generate immunity or how long antibodies will persist. Lab results should not be used as the sole basis to diagnose or exclude recent SARS-CoV-2 infection. Correlation with epidemiologic risk factors and other clinical and laboratory findings is recommended.